

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Mucin 5AC (MUC5AC)(MUC5AC/917), CF568 conjugate, 0.1mg/mL, Clone: [MUC5AC/917], Mouse, Monoclonal BOT-BNC680917-100**

Artikelname	Mucin 5AC (MUC5AC)(MUC5AC/917), CF568 conjugate, 0.1mg/mL, Clone: [MUC5AC/917], Mouse, Monoclonal
Artikelnummer	BOT-BNC680917-100
Hersteller Artikelnummer	BNC680917-100
Alternativnummer	BOT-BNC680917-100-100UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Human
Immunogen	Recombinant human MUC5AC protein
Konjugation	CF568
Produktbeschreibung	Mucin 5AC glycoprotein (MUC5AC) is a 641 kDa glycoprotein belonging to the superfamily of mucins. Mucins are high molecular weight glycoproteins produced by epithelial cells and can be divided into two families, secretory mucins and membrane bound mu...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL
Klon-Bezeichnung	[MUC5AC/917]

Molekulargewicht	641 kDa
UniProt	<a href="#">P98088</a>
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology formalin-fixed 1-2 ug/mL Does not react with rat, others not known Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user