

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

von Willebrand Factor / Factor VIII Related-Ag (Endothelial Marker)(IIIIE2.34), CF568 conjugate, 0.1mg/mL, Clone: [IIIIE2.34], Mouse, Monoclonal BOT-BNC680934-500

Artikelname	von Willebrand Factor / Factor VIII Related-Ag (Endothelial Marker)(IIIIE2.34), CF568 conjugate, 0.1mg/mL, Clone: [IIIIE2.34], Mouse, Monoclonal
Artikelnummer	BOT-BNC680934-500
Hersteller Artikelnummer	BNC680934-500
Alternativnummer	BOT-BNC680934-500-500UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Human
Immunogen	Recombinant human vWF fragment spanning aa 845-949
Konjugation	CF568
Produktbeschreibung	von Willebrand Factor (vWF) is a multimeric glycoprotein that is found in endothelial cells, plasma and platelets. It acts as a carrier protein for Factor VIII and promotes platelet adhesion and aggregation. vWF undergoes a variety of posttranslation...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL

Klon-Bezeichnung	[IIIIE2.34]
Molekulargewicht	250 kDa
UniProt	P04275
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistochemistry (formalin-fixed): 0.5-1.0 ug/mL for 30 minutes at RT Immunoprecipitation: 0.5-1 ug/500 ug protein lysate Western blot: 0.5-1.0 ug/mL Flow cytometry: 0.5-1 ug/million cells Immunofluorescence: 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user