

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Cytokeratin 5/8(C-50), CF568 conjugate, 0.1mg/mL, Clone: [C-50], Mouse, Monoclonal BOT-BNC680948-100**

|                          |   |
|--------------------------|---|
| Artikelname              | Cytokeratin 5/8(C-50), CF568 conjugate, 0.1mg/mL, Clone: [C-50], Mouse, Monoclonal  |
| Artikelnummer            | BOT-BNC680948-100   |
| Hersteller Artikelnummer | BNC680948-100   |
| Alternativnummer         | BOT-BNC680948-100-100UL   |
| Hersteller               | Biotium   |
| Wirt                     | Mouse   |
| Kategorie                | Antikörper  |
| Applikation              | IHC   |
| Spezies Reaktivität      | Bovine, Canine, Ferret, Hamster, Human, Mouse, Porcine, Rat, Sheep  |
| Immunogen                | Cytoskeletal preparation from HeLa cells  |
| Konjugation              | CF568   |
| Produktbeschreibung      | This antibody reacts with keratin 5 (58 kDa) and keratin 8 (52.5 kDa). Simple epithelia express cytokeratin 8 in combination with 18. On the other hand, basal cells of stratified epithelia express cytokeratin 5 paired with 14. This antibody therefore... |
| Klonalität               | Monoclonal  |
| Konzentration            | 0.1 mg/mL   |
| Klon-Bezeichnung         | [C-50]  |

|                        |  |
|------------------------|--|
| Molekulargewicht       | 58 kDa (CK5), 52.5 kDa (CK8)   |
| UniProt                | <a href="#">P13647</a>   |
| Puffer                 | PBS, 0.1% BSA, 0.05% azide   |
| Quelle                 | Animal   |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology formalin-fixed 0.5-1.0 ug/mL For staining of formalin-fixed tissues, digest sections with trypsin at 1mg/mL PBS, 15 min at RT Flow Cytometry 0.5-1 ug/million cells/0.1 mL Immunofluorescence 0.5-1.0 ug/mL Western blotting 0.5-1.0 ug/mL Does not react with chicken , or Xenopus Optimal dilution for a specific application should be determined by user |