

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Mucin 1 / EMA / Episialin / CD227(MUC1/955), CF568 conjugate, 0.1mg/mL, Clone: [MUC1/955], Mouse, Monoclonal BOT-BNC680955-500

| | |
|--------------------------|---|
| Artikelname | Mucin 1 / EMA / Episialin / CD227(MUC1/955), CF568 conjugate, 0.1mg/mL, Clone: [MUC1/955], Mouse, Monoclonal |
| Artikelnummer | BOT-BNC680955-500 |
| Hersteller Artikelnummer | BNC680955-500 |
| Alternativnummer | BOT-BNC680955-500-500UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC, WB |
| Spezies Reaktivität | Human, Mouse |
| Immunogen | Human milk-fat globule membranes (HMFGM) |
| Konjugation | CF568 |
| Produktbeschreibung | This MAb reacts with MUC1. The dominant epitope of this MAb has not yet been determined. MUC1 is a large cell surface mucin glycoprotein expressed by most glandular and ductal epithelial cells and some hematopoietic cell lineages. It is expressed on ... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |
| Klon-Bezeichnung | [MUC1/955] |

| | |
|------------------------|---|
| Molekulargewicht | 265-400 kDa |
| UniProt | P15941 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology formalin-fixed 0.25-0.5 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL ELISA For coating use Ab at 2-5 ug/mL, order Ab without BSA Optimal dilution for a specific application should be determined by user |