

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Mucin 1 / EMA / Episialin / CD227(MUC1/967), CF568 conjugate, 0.1mg/mL, Clone: [MUC1/967], Mouse, Monoclonal BOT-BNC680967-500**

Artikelname	Mucin 1 / EMA / Episialin / CD227(MUC1/967), CF568 conjugate, 0.1mg/mL, Clone: [MUC1/967], Mouse, Monoclonal
Artikelnummer	BOT-BNC680967-500
Hersteller Artikelnummer	BNC680967-500
Alternativnummer	BOT-BNC680967-500-500UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC, WB
Spezies Reaktivität	Human
Immunogen	Human milk-fat globule membranes (HMFGM)
Konjugation	CF568
Produktbeschreibung	This MAb recognizes full-length MUC1 in a glycosylation-independent manner and can bind to the fully glycosylated protein. The dominant epitope of this MAb is APDTR in the VNTR region. It reacts with the core peptide of the MUC1 protein, which is a m...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL
Klon-Bezeichnung	[MUC1/967]

Molekulargewicht	265-400 kDa
UniProt	<a href="#">P15941</a>
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Immunohistology formalin-paraffin 0.25-0.5 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Immunofluorescence 1-2 ug/mL Optimal dilution for a specific application should be determined by user