

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Cytokeratin 8/18(KRT8/803 + KRT18/835), CF568 conjugate, 0.1mg/mL, Clone: [KRT8/803 KRT18/835], Mouse, Monoclonal BOT-BNC680979-500

Artikelname	Cytokeratin 8/18(KRT8/803 + KRT18/835), CF568 conjugate, 0.1mg/mL, Clone: [KRT8/803 KRT18/835], Mouse, Monoclonal
Artikelnummer	BOT-BNC680979-500
Hersteller Artikelnummer	BNC680979-500
Alternativnummer	BOT-BNC680979-500-500UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Human
Immunogen	Recombinant full-length human KRT8 protein (KRT8/803), Recombinant full-length human KRT18 protein (KRT18/835)
Konjugation	CF568
Produktbeschreibung	Cytokeratin 8 (CK8) belongs to the type II (or B or basic) subfamily of high molecular weight cytokeratins and exists in combination with cytokeratin 18 (CK18). This MAbs cocktail recognizes all simple epithelia including glandular epithelium, for exa...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL

Klon-Bezeichnung	[KRT8/803 KRT18/835]
Molekulargewicht	52.5 kDa (CK8), 45 kDa (CK18)
UniProt	P05787
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	<p>Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology formalin-fixed 0.5-1.0 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Western blotting 0.5-1.0 ug/mL Optimal dilution for a specific application should be determined by user</p>