

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Cytokeratin, pan (Cocktail PAN-CK), CF568 conjugate, 0.1mg/mL, Clone: [Cocktail PAN-CK], Mouse, Monoclonal BOT-BNC680999-100**

|                          |   |
|--------------------------|---|
| Artikelname              | Cytokeratin, pan (Cocktail PAN-CK), CF568 conjugate, 0.1mg/mL, Clone: [Cocktail PAN-CK], Mouse, Monoclonal  |
| Artikelnummer            | BOT-BNC680999-100   |
| Hersteller Artikelnummer | BNC680999-100   |
| Alternativnummer         | BOT-BNC680999-100-100UL   |
| Hersteller               | Biotium   |
| Wirt                     | Mouse   |
| Kategorie                | Antikörper  |
| Applikation              | FC, IF, IHC   |
| Spezies Reaktivität      | Bovine, Canine, Gallus, Human, Monkey, Mouse, Rabbit, Rat   |
| Immunogen                | Human epidermal keratin   |
| Konjugation              | CF568   |
| Produktbeschreibung      | Twenty human keratins are resolved with two-dimensional gel electrophoresis into acidic (pI 6.0) subfamilies. This antibody cocktail recognizes acidic (Type I or LMW) and basic (Type II or HMW) cytokeratins, with 67 kDa (CK1), 64 kDa (CK3), 59 kDa (C... |
| Klonalität               | Monoclonal  |
| Konzentration            | 0.1 mg/mL   |
| Klon-Bezeichnung         | [Cocktail PAN-CK]   |

|                        |   |
|------------------------|---|
| Molekulargewicht       | 40-67 kDa (Multiple)  |
| UniProt                | <a href="#">Q7Z794</a>  |
| Puffer                 | PBS, 0.1% BSA, 0.05% azide  |
| Quelle                 | Animal  |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology formalin-fixed 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Western blotting 0.5-1 ug/mL Optimal dilution for a specific application should be determined by user |