

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

CD11c (Dendritic Cell Marker) (ITGAX/1284), CF568 conjugate, 0.1mg/mL, IgG2b, Clone: [ITGAX/1284], Mouse, Monoclonal BOT-BNC681284-500

| | |
|--------------------------|---|
| Artikelname | CD11c (Dendritic Cell Marker) (ITGAX/1284), CF568 conjugate, 0.1mg/mL, IgG2b, Clone: [ITGAX/1284], Mouse, Monoclonal |
| Artikelnummer | BOT-BNC681284-500 |
| Hersteller Artikelnummer | BNC681284-500 |
| Alternativnummer | BOT-BNC681284-500-500UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant human ITGAX protein fragment (aa 637-827) (exact sequence is proprietary) |
| Konjugation | CF568 |
| Produktbeschreibung | This antibody recognizes a protein of 145 kDa, identified as CD11c. CD11c (ITGAX), a member of the leukointegrin family, shares the same beta subunit with other members of the leukocyte adhesion molecule family, which includes CD11a (LFA-1), CD11b (M... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |

| | |
|------------------------|---|
| Klon-Bezeichnung | [ITGAX/1284] |
| Molekulargewicht | 145 kDa |
| Isotyp | IgG2b |
| UniProt | P20702 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology (formalin) 1-2 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM Tris, 1 mM EDTA pH 9.0 for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user |