

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Erythropoietin (EPO) (EPO/1368), CF568 conjugate, 0.1mg/mL, Clone: [EPO/1368], Mouse, Monoclonal BOT-BNC681368-500

| | |
|--------------------------|---|
| Artikelname | Erythropoietin (EPO) (EPO/1368), CF568 conjugate, 0.1mg/mL, Clone: [EPO/1368], Mouse, Monoclonal |
| Artikelnummer | BOT-BNC681368-500 |
| Hersteller Artikelnummer | BNC681368-500 |
| Alternativnummer | BOT-BNC681368-500-500UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC, WB |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant fragment of human EPO protein (aa28-162) (exact sequence is proprietary) |
| Konjugation | CF568 |
| Produktbeschreibung | This antibody recognizes a protein of about 37 kDa, which is identified as Erythropoietin (EPO). Erythropoietin is a secreted, glycosylated cytokine hormone composed of four alpha helical bundles. It is the primary factor responsible for regulating e... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |

| | |
|------------------------|--|
| Klon-Bezeichnung | [EPO/1368] |
| Molekulargewicht | 37 kDa |
| UniProt | P01588 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | <p>Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology (formalin) 4-8 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL Western blotting 0.5-1 ug/mL Optimal dilution for a specific application should be determined by user</p> |