

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Wilms Tumor 1 (WT1) (Wilms Tumor & Mesothelial Marker) (WT1/1434R), CF568 conjugate, 0.1mg/mL, Clone: [WT1/1434R], Rabbit, Monoclonal BOT-BNC681434-500

| | |
|--------------------------|---|
| Artikelname | Wilms Tumor 1 (WT1) (Wilms Tumor & Mesothelial Marker) (WT1/1434R), CF568 conjugate, 0.1mg/mL, Clone: [WT1/1434R], Rabbit, Monoclonal |
| Artikelnummer | BOT-BNC681434-500 |
| Hersteller Artikelnummer | BNC681434-500 |
| Alternativnummer | BOT-BNC681434-500-500UL |
| Hersteller | Biotium |
| Wirt | Rabbit |
| Kategorie | Antikörper |
| Applikation | IHC |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant human full-length WT1 protein |
| Konjugation | CF568 |
| Produktbeschreibung | This antibody recognizes a 47-55 kDa-tumor suppressor protein, identified as Wilms Tumor (WT1) protein. The antibody reacts with all isoforms of the full-length WT1 and also identifies WT1 lacking exon 2-encoded amino acids, frequently found in subse... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |

| | |
|------------------------|--|
| Klon-Bezeichnung | [WT1/1434R] |
| Molekulargewicht | 47-55 kDa |
| UniProt | P19544 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | <p>Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology (formalin): 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user</p> |