

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Growth Hormone (Pituitary Marker) (GH/1450), CF568 conjugate, 0.1mg/mL, IgG2b, Clone: [GH/1450], Mouse, Monoclonal BOT-BNC681450-100

| | |
|--------------------------|---|
| Artikelname | Growth Hormone (Pituitary Marker) (GH/1450), CF568 conjugate, 0.1mg/mL, IgG2b, Clone: [GH/1450], Mouse, Monoclonal |
| Artikelnummer | BOT-BNC681450-100 |
| Hersteller Artikelnummer | BNC681450-100 |
| Alternativnummer | BOT-BNC681450-100-100UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC |
| Spezies Reaktivität | Human |
| Immunogen | A recombinant human Growth Hormone (GH) fragment (aa58-187) (exact sequence is proprietary) |
| Konjugation | CF568 |
| Produktbeschreibung | Pituitary growth hormone (GH) plays a crucial role in stimulating and controlling the growth, metabolism and differentiation of many mammalian cell types by modulating the synthesis of multiple mRNA species. These effects are mediated by the binding ... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |

| | |
|------------------------|---|
| Klon-Bezeichnung | [GH/1450] |
| Molekulargewicht | 20 kDa |
| Isotyp | IgG2b |
| UniProt | P01241 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology (formalin): 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user |