

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Alpha-1-Antichymotrypsin (SERPINA3) (Histiocytoma Marker) (AACT/1451 + AACT/1452), CF568 conjugate, 0.1mg/mL, IgG1, Clone: [AACT/1451 AACT/1452], Mouse, Monoclonal BOT-BNC681453-500**

Artikelname	Alpha-1-Antichymotrypsin (SERPINA3) (Histiocytoma Marker) (AACT/1451 + AACT/1452), CF568 conjugate, 0.1mg/mL, IgG1, Clone: [AACT/1451 AACT/1452], Mouse, Monoclonal
Artikelnummer	BOT-BNC681453-500
Hersteller Artikelnummer	BNC681453-500
Alternativnummer	BOT-BNC681453-500-500UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Human
Immunogen	Recombinant human Antichymotrypsin (AACT) protein fragment (aa49-187) (exact sequence is proprietary)
Konjugation	CF568
Produktbeschreibung	Alpha-1 Antichymotrypsin (AACT) is a plasma protease inhibitor synthesized in the liver as a single glycopeptide chain. In human, the normal serum level of AACT is about one-tenth that of alpha-1-antitrypsin (AAT), with which it shares nucleic acid a...
Klonalität	Monoclonal

Konzentration	0.1 mg/mL
Klon-Bezeichnung	[AACT/1451 AACT/1452]
Molekulargewicht	65-76 kDa
Isotyp	IgG1
UniProt	<a href="#">P01011</a>
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Immunofluorescence 0.5-1 ug/mL Western blotting 0.5-1 ug/mL Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user