

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **FOXA1 / HNF3A (FOXA1/1519), CF568 conjugate, 0.1mg/mL, IgG2a, Clone: [FOXA1/1519], Mouse, Monoclonal BOT-BNC681519-100**

|                          |   |
|--------------------------|---|
| Artikelname              | FOXA1 / HNF3A (FOXA1/1519), CF568 conjugate, 0.1mg/mL, IgG2a, Clone: [FOXA1/1519], Mouse, Monoclonal  |
| Artikelnummer            | BOT-BNC681519-100   |
| Hersteller Artikelnummer | BNC681519-100   |
| Alternativnummer         | BOT-BNC681519-100-100UL   |
| Hersteller               | Biotium   |
| Wirt                     | Mouse   |
| Kategorie                | Antikörper  |
| Applikation              | FC, IHC, WB   |
| Spezies Reaktivität      | Human, Rat  |
| Immunogen                | Recombinant human FOXA1 protein fragment (aa467-614) (exact sequence is proprietary)  |
| Konjugation              | CF568   |
| Produktbeschreibung      | The transcription factor Forkhead-box A1 (FOXA1), also known as hepatocyte nuclear factor 3-alpha, is a member of the FOX class of transcription factors. HNF-1 (alpha and beta), HNF-3 (alpha, beta and gamma), HNF-4 (alpha and gamma), and HNF-6 compos... |
| Klonalität               | Monoclonal  |
| Konzentration            | 0.1 mg/mL   |

|                        |   |
|------------------------|---|
| Klon-Bezeichnung       | [FOXA1/1519]  |
| Molekulargewicht       | 79 kDa  |
| Isotyp                 | IgG2a   |
| UniProt                | <a href="#">P55317</a>  |
| Puffer                 | PBS, 0.1% BSA, 0.05% azide  |
| Quelle                 | Animal  |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology (formalin) 1-2 ug/mL Staining of formalin-fixed tissues is enhanced by boiling tissue sections in 10 mM Tris, 1 mM EDTA pH 9.0 for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user |