

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Cytokeratin, Basic (Type II or HMW) (Epithelial Marker) (KRTH/1576R), CF568 conjugate, 0.1mg/mL, Clone: [KRTH/1576R], Rabbit, Monoclonal BOT-BNC681576-100

Artikelname	Cytokeratin, Basic (Type II or HMW) (Epithelial Marker) (KRTH/1576R), CF568 conjugate, 0.1mg/mL, Clone: [KRTH/1576R], Rabbit, Monoclonal
Artikelnummer	BOT-BNC681576-100
Hersteller Artikelnummer	BNC681576-100
Alternativnummer	BOT-BNC681576-100-100UL
Hersteller	Biotium
Wirt	Rabbit
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Human
Immunogen	Recombinant full-length human KRT76 protein
Konjugation	CF568
Produktbeschreibung	This MAb recognizes basic (Type II or HMW) cytokeratins, which include 67 kDa (CK1), 64 kDa (CK3), 59 kDa (CK4), 58 kDa (CK5), 56 kDa (CK6), 52 kDa (CK8). Twenty human keratins are resolved with two-dimensional gel electrophoresis into acidic (pI 6.0...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL

Klon-Bezeichnung	[KRTH/1576R]
Molekulargewicht	52-67 kDa
UniProt	Q01546
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 0.25-0.5 ug/mL Immunofluorescence 1-2 ug/mL Western blotting 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user