

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

CFTR (Cystic Fibrosis Transmembrane Conductance Regulator) (CFTR/1785), CF568 conjugate, 0.1mg/mL, IgG2b, Clone: [CFTR/1785], Mouse, Monoclonal BOT-BNC681785-100

Artikelname	CFTR (Cystic Fibrosis Transmembrane Conductance Regulator) (CFTR/1785), CF568 conjugate, 0.1mg/mL, IgG2b, Clone: [CFTR/1785], Mouse, Monoclonal
Artikelnummer	BOT-BNC681785-100
Hersteller Artikelnummer	BNC681785-100
Alternativnummer	BOT-BNC681785-100-100UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Human
Immunogen	Recombinant human CFTR fragment (aa258-385) (exact sequence is proprietary)
Konjugation	CF568
Produktbeschreibung	This antibody recognizes a protein of 165-170 kDa, identified as cystic fibrosis transmembrane conductance regulator (CFTR). CFTR is composed of two membrane-spanning domains (MSD), two nucleotide-binding domains (NBD), and an R domain. It is structu...
Klonalität	Monoclonal

Konzentration	0.1 mg/mL
Klon-Bezeichnung	[CFTR/1785]
Molekulargewicht	165-170 kDa
Isotyp	IgG2b
UniProt	P13569
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology (formalin) 1-2 ug/mL Staining of formalin-fixed tissues is enhanced by boiling tissue sections in 10 mM Tris, 1 mM EDTA pH 9.0 for 10-20 min followed by cooling at RT for 20 min Optimal dilution for a specific application should be determined by user