

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

CD20 / MS4A1 (B-Cell Marker) (rIGEL/773), CF568 conjugate, 0.1mg/mL, Clone: [rIGEL/773], Mouse, Monoclonal BOT-BNC681829-500

Artikelname	CD20 / MS4A1 (B-Cell Marker) (rIGEL/773), CF568 conjugate, 0.1mg/mL, Clone: [rIGEL/773], Mouse, Monoclonal
Artikelnummer	BOT-BNC681829-500
Hersteller Artikelnummer	BNC681829-500
Alternativnummer	BOT-BNC681829-500-500UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	FC, IF, IHC
Spezies Reaktivität	Human
Immunogen	Recombinant full-length human MS4A1 protein
Konjugation	CF568
Produktbeschreibung	This antibody recognizes a protein of 30-33 kDa, which is identified as CD20. It is a non-Ig differentiation antigen of B-cells and its expression is restricted to normal and neoplastic B-cells, being absent from all other leukocytes and tissues. CD2...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL
Klon-Bezeichnung	[rIGEL/773]

Molekulargewicht	33-37 kDa
UniProt	P11836
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology (formalin): 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user