

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Emerin (Papillary Thyroid Carcinoma and EDMD Marker) (EMD/2167), CF568 conjugate, 0.1mg/mL, IgG1, Clone: [EMD/2167], Mouse, Monoclonal BOT-BNC682167-100

| | |
|--------------------------|---|
| Artikelname | Emerin (Papillary Thyroid Carcinoma and EDMD Marker) (EMD/2167), CF568 conjugate, 0.1mg/mL, IgG1, Clone: [EMD/2167], Mouse, Monoclonal |
| Artikelnummer | BOT-BNC682167-100 |
| Hersteller Artikelnummer | BNC682167-100 |
| Alternativnummer | BOT-BNC682167-100-100UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IF, IHC, WB |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant human Emerin protein fragment (around aa 56-167) (exact sequence is proprietary) |
| Konjugation | CF568 |
| Produktbeschreibung | Emerin is a member of the nuclear lamina associated protein family. It is ubiquitously expressed and localized to the nuclear membrane in normal cells. Mutations of the gene that encodes emerin result in the X-linked recessive disease Emery-Dreifuss ... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |

| | |
|------------------------|---|
| Klon-Bezeichnung | [EMD/2167] |
| Molekulargewicht | 37 kDa |
| Isotyp | IgG1 |
| UniProt | P50402 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 1-2 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user |