

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Catenin, beta (CTNNB1) (rCTNNB1/2173), CF568 conjugate, 0.1mg/mL, Clone: [rCTNNB1/2173], Mouse, Monoclonal BOT-BNC682173-100**

Artikelname	Catenin, beta (CTNNB1) (rCTNNB1/2173), CF568 conjugate, 0.1mg/mL, Clone: [rCTNNB1/2173], Mouse, Monoclonal
Artikelnummer	BOT-BNC682173-100
Hersteller Artikelnummer	BNC682173-100
Alternativnummer	BOT-BNC682173-100-100UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IF, IHC, WB
Spezies Reaktivität	Human, Mouse, Rat
Immunogen	Recombinant full-length human i-catenin protein
Konjugation	CF568
Produktbeschreibung	Beta-catenin associates with the cytoplasmic portion of E-cadherin. The catenin/cadherin complexes play an important role mediating cellular adhesion, including adherens junctions. Beta-catenin is involved in the Wnt signaling pathway as well as othe...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL
Klon-Bezeichnung	[rCTNNB1/2173]

Molekulargewicht	92 kDa
UniProt	<a href="#">P35222</a>
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology (formalin): 1-2 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM Tris with 1 mM EDTA pH 9.0 for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user