

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

E-Cadherin (CDH1) / CD324 (Intercellular Junction Marker) (CDH1/2208R), CF568 conjugate, 0.1mg/mL, Clone: [CDH1/2208R], Rabbit, Monoclonal BOT-BNC682208-500

| | |
|--------------------------|---|
| Artikelname | E-Cadherin (CDH1) / CD324 (Intercellular Junction Marker) (CDH1/2208R), CF568 conjugate, 0.1mg/mL, Clone: [CDH1/2208R], Rabbit, Monoclonal |
| Artikelnummer | BOT-BNC682208-500 |
| Hersteller Artikelnummer | BNC682208-500 |
| Alternativnummer | BOT-BNC682208-500-500UL |
| Hersteller | Biotium |
| Wirt | Rabbit |
| Kategorie | Antikörper |
| Applikation | FC, IF, IHC, WB |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant full-length human E-Cadherin protein |
| Konjugation | CF568 |
| Produktbeschreibung | This antibody recognizes a protein of 120-80 kDa, identified as E-cadherin. Cadherins comprise a family of calcium-dependent adhesion molecules that function to mediate cell-cell binding critical to the maintenance of tissue structure and morphogenes... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |

| | |
|------------------------|--|
| Klon-Bezeichnung | [CDH1/2208R] |
| Molekulargewicht | 120-80 kDa (Mature), 135 kDa (Precursor) |
| UniProt | P12830 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Does not react with mouse or rat Immunohistology (formalin): 1-2 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues is enhanced by boiling tissue sections in 10 mM Tris with 1 mM EDTA pH 9.0 for 10-20 minutes followed by cooling at RT for 20 minutes Western blotting 0.5-1 ug/mL Optimal dilution for a specific application should be determined by user |