

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

CD43 (T-Cell Marker) (rSPN/839), CF568 conjugate, 0.1mg/mL, IgG1, Clone: [rSPN/839], Mouse, Monoclonal BOT-BNC682220-100

| | |
|--------------------------|---|
| Artikelname | CD43 (T-Cell Marker) (rSPN/839), CF568 conjugate, 0.1mg/mL, IgG1, Clone: [rSPN/839], Mouse, Monoclonal |
| Artikelnummer | BOT-BNC682220-100 |
| Hersteller Artikelnummer | BNC682220-100 |
| Alternativnummer | BOT-BNC682220-100-100UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | FC, IF, IHC, WB |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant full-length human CD43 protein |
| Konjugation | CF568 |
| Produktbeschreibung | This antibody recognizes a cell surface glycoprotein of 95/115/135 kDa (depending upon the extent of glycosylation), identified as CD43 . Epitope of MAb Bra7G is clearly different from that of MAb rSPN/839, called b as opposed to a for rSPN/839. 70-9... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |
| Klon-Bezeichnung | [rSPN/839] |

| | |
|------------------------|--|
| Molekulargewicht | 95, 115, or 135 kDa |
| Isotyp | IgG1 |
| UniProt | P16150 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | <p>Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 0.5-1 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user</p> |