

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Bcl-2 (Apoptosis & Follicular Lymphoma Marker) (rBCL2/782), CF568 conjugate, 0.1mg/mL, Clone: [rBCL2/782], Mouse, Monoclonal BOT-BNC682221-500

Artikelname	Bcl-2 (Apoptosis & Follicular Lymphoma Marker) (rBCL2/782), CF568 conjugate, 0.1mg/mL, Clone: [rBCL2/782], Mouse, Monoclonal
Artikelnummer	BOT-BNC682221-500
Hersteller Artikelnummer	BNC682221-500
Alternativnummer	BOT-BNC682221-500-500UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC, WB
Spezies Reaktivität	Human
Immunogen	Recombinant full-length human BCL2 protein
Konjugation	CF568
Produktbeschreibung	This antibody recognizes a protein of 25-26 kDa, identified as Bcl-2. This antibody shows no cross-reaction with Bcl-x or Bax proteins. Bcl-2 suppresses apoptosis in a variety of cell systems including factor-dependent lymphohematopoietic and neural ...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL
Klon-Bezeichnung	[rBCL2/782]

Molekulargewicht	25-26 kDa
UniProt	P10415
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 0.5-1 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues is enhanced by boiling tissue sections in 1 mM EDTA Buffer pH 8.0 for 10-20 minutes followed by cooling at RT for 20 minutes Western blotting 0.5-1 ug/mL Optimal dilution for a specific application should be determined by user