

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Villin (GI-Mucosal & Urogenital Brush Border Marker) (rVIL1/1325), CF568 conjugate, 0.1mg/mL, Clone: [rVIL1/1325], Mouse, Monoclonal BOT-BNC682279-100**

Artikelname	Villin (GI-Mucosal & Urogenital Brush Border Marker) (rVIL1/1325), CF568 conjugate, 0.1mg/mL, Clone: [rVIL1/1325], Mouse, Monoclonal
Artikelnummer	BOT-BNC682279-100
Hersteller Artikelnummer	BNC682279-100
Alternativnummer	BOT-BNC682279-100-100UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Human
Immunogen	Recombinant human Villin fragment of 133 amino acid residues (aa179-311)
Konjugation	CF568
Produktbeschreibung	This antibody recognizes a protein of 95 kDa, which is identified as villin. It is a major constituent in the microvilli, which compose the brush border of epithelial cells forming absorptive surfaces of the intestinal and renal proximal tubular epit...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL

Klon-Bezeichnung	[rVIL1/1325]
Molekulargewicht	93 kDa
UniProt	<a href="#">P09327</a>
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 1-2 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user