

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Estrogen Receptor, alpha (Marker of Estrogen Dependence) (rESR1/1935), CF568 conjugate, 0.1mg/mL, IgG1, Clone: [rESR1/1935], Mouse, Monoclonal BOT-BNC682300-100**

Artikelname	Estrogen Receptor, alpha (Marker of Estrogen Dependence) (rESR1/1935), CF568 conjugate, 0.1mg/mL, IgG1, Clone: [rESR1/1935], Mouse, Monoclonal
Artikelnummer	BOT-BNC682300-100
Hersteller Artikelnummer	BNC682300-100
Alternativnummer	BOT-BNC682300-100-100UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC, WB
Spezies Reaktivität	Human
Immunogen	Recombinant full-length human Estrogen Receptor alpha protein
Konjugation	CF568
Produktbeschreibung	This MAb is specific to ER alpha and shows minimal cross-reaction with other members of the family. ER is an important regulator of growth and differentiation in the mammary gland. Presence of ER in breast tumors indicates an increased likelihood of ...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL

Klon-Bezeichnung	[rESR1/1935]
Molekulargewicht	67 kDa
Isotyp	IgG1
UniProt	<a href="#">P03372</a>
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 1-2 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Western blotting 1-2 ug/mL Optimal dilution for a specific application should be determined by user