

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **CD11c (Dendritic Cell Marker)(ITGAX/1242), CF594 conjugate, 0.1mg/mL, Clone: [ITGAX/1242], Mouse, Monoclonal BOT-BNC941242-100**

|                          |   |
|--------------------------|---|
| Artikelname              | CD11c (Dendritic Cell Marker)(ITGAX/1242), CF594 conjugate, 0.1mg/mL, Clone: [ITGAX/1242], Mouse, Monoclonal  |
| Artikelnummer            | BOT-BNC941242-100   |
| Hersteller Artikelnummer | BNC941242-100   |
| Alternativnummer         | BOT-BNC941242-100-100UL   |
| Hersteller               | Biotium   |
| Wirt                     | Mouse   |
| Kategorie                | Antikörper  |
| Applikation              | IHC   |
| Spezies Reaktivität      | Human   |
| Immunogen                | Recombinant human ITGAX protein fragment (aa 637-827) (exact sequence is proprietary)   |
| Konjugation              | CF594   |
| Produktbeschreibung      | Recognizes a protein of 145 kDa, identified as CD11c. CD11c (ITGAX), a member of the leukointegrin family, shares the same beta subunit with other members of the leukocyte adhesion molecule family, which includes CD11a (LFA-1), CD11b (MAC-1) and CD11... |
| Klonalität               | Monoclonal  |
| Konzentration            | 0.1 mg/mL   |

|                        |  |
|------------------------|--|
| Klon-Bezeichnung       | [ITGAX/1242]   |
| Molekulargewicht       | 145 kDa  |
| UniProt                | <a href="#">P20702</a>   |
| Puffer                 | PBS, 0.1% BSA, 0.05% azide   |
| Quelle                 | Animal   |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistochemistry (formalin-fixed): 1-2 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user |