

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Interleukin 10 (IL10) (Immunoregulation Marker) (IL10/2651R), CF594 conjugate, 0.1mg/mL, Clone: [IL10/2651R], Rabbit, Monoclonal BOT-BNC942651-100

| | |
|--------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Artikelname | Interleukin 10 (IL10) (Immunoregulation Marker) (IL10/2651R), CF594 conjugate, 0.1mg/mL, Clone: [IL10/2651R], Rabbit, Monoclonal |
| Artikelnummer | BOT-BNC942651-100 |
| Hersteller Artikelnummer | BNC942651-100 |
| Alternativnummer | BOT-BNC942651-100-100UL |
| Hersteller | Biotium |
| Wirt | Rabbit |
| Kategorie | Antikörper |
| Applikation | IHC |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant full-length human Interleukin 10 (IL10) protein |
| Konjugation | CF594 |
| Produktbeschreibung | Interleukin 10 (IL10) is a cytokine produced primarily by monocytes and to a lesser extent by lymphocytes. This cytokine has pleiotropic effects in immuno-regulation and inflammation. It down-regulates the expression of Th1 cytokines, MHC class II an... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |

| | |
|------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Klon-Bezeichnung | [IL10/2651R] |
| Molekulargewicht | 20 kDa (Monomer), 37 kDa (Dimer) |
| UniProt | P22301 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 1-2 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user |