

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Interleukin 10 (IL10) (Immunoregulation Marker) (IL10/2651R), CF594 conjugate, 0.1mg/mL, Clone: [IL10/2651R], Rabbit, Monoclonal BOT-BNC942651-500**

Artikelname	Interleukin 10 (IL10) (Immunoregulation Marker) (IL10/2651R), CF594 conjugate, 0.1mg/mL, Clone: [IL10/2651R], Rabbit, Monoclonal
Artikelnummer	BOT-BNC942651-500
Hersteller Artikelnummer	BNC942651-500
Alternativnummer	BOT-BNC942651-500-500UL
Hersteller	Biotium
Wirt	Rabbit
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Human
Immunogen	Recombinant full-length human Interleukin 10 (IL10) protein
Konjugation	CF594
Produktbeschreibung	Interleukin 10 (IL10) is a cytokine produced primarily by monocytes and to a lesser extent by lymphocytes. This cytokine has pleiotropic effects in immuno-regulation and inflammation. It down-regulates the expression of Th1 cytokines, MHC class II an...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL

Klon-Bezeichnung	[IL10/2651R]
Molekulargewicht	20 kDa (Monomer), 37 kDa (Dimer)
UniProt	<a href="#">P22301</a>
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 1-2 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user