

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

B7-H4 (Immuno-Inhibitory Protein) (B7H4/2652R), CF594 conjugate, 0.1mg/mL, Clone: [B7H4/2652R], Rabbit, Monoclonal BOT-BNC942652-500

Artikelname	B7-H4 (Immuno-Inhibitory Protein) (B7H4/2652R), CF594 conjugate, 0.1mg/mL, Clone: [B7H4/2652R], Rabbit, Monoclonal
Artikelnummer	BOT-BNC942652-500
Hersteller Artikelnummer	BNC942652-500
Alternativnummer	BOT-BNC942652-500-500UL
Hersteller	Biotium
Wirt	Rabbit
Kategorie	Antikörper
Applikation	FC, IF, IHC
Spezies Reaktivität	Human
Immunogen	Recombinant human B7-H4 protein fragment (exact sequence is proprietary)
Konjugation	CF594
Produktbeschreibung	B7-H4 (B7-homolog 4), belongs to the B7 immunoglobulin superfamily of ligand-lymphocyte interacting proteins. Expressed primarily on the membrane of lymphoid cells, B7-H4 is an immuno-inhibitory protein that interacts with receptors on the surface of...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL

Klon-Bezeichnung	[B7H4/2652R]
Molekulargewicht	35 kDa
UniProt	Q7Z7D3
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody ELISA: 2-4 ug/m for coating order Ab without BSA Immunohistology (formalin): 1-2 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user