

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

HL-60 Antigen / Human Leukemia Antigen (IPO-M6), Biotin conjugate, 0.1mg/mL, IgG1, Clone: [IPO-M6], Mouse, Monoclonal BOT-BNCB0228-100

| | |
|--------------------------|---|
| Artikelname | HL-60 Antigen / Human Leukemia Antigen (IPO-M6), Biotin conjugate, 0.1mg/mL, IgG1, Clone: [IPO-M6], Mouse, Monoclonal |
| Artikelnummer | BOT-BNCB0228-100 |
| Hersteller Artikelnummer | BNCB0228-100 |
| Alternativnummer | BOT-BNCB0228-100-100UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Spezies Reaktivität | Human |
| Immunogen | Leukemia HL-60 cells |
| Konjugation | Biotin |
| Produktbeschreibung | This antibody reacts with human leukemia cell line HL-60 and immunoprecipitates two proteins with MW of 48 kDa and 52 kDa. It does not stain B-cell lines Daudi, PHS, Namalwa, RPMI-1788 and T-cell lines CCRF-HSB2, Jurkat and Molt-4. This antibody stai... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |
| Klon-Bezeichnung | [IPO-M6] |
| Molekulargewicht | 48 kDa & 52 kDa |

| | |
|------------------------|--|
| Isotyp | IgG1 |
| UniProt | Not Known |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | For coating for ELISA, order Ab without BSA Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Optimal dilution and staining procedure for a specific application should be determined by user Recommended starting concentrations for titration are 1-2 ug/mL for most applications, or 1 ug/million cells/100 uL for flow cytometry |