

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Cytokeratin, LMW(AE-1), Biotin conjugate, 0.1mg/mL, Clone: [AE-1], Mouse, Monoclonal BOT-BNCB0253-500

| | |
|--------------------------|---|
| Artikelname | Cytokeratin, LMW(AE-1), Biotin conjugate, 0.1mg/mL, Clone: [AE-1], Mouse, Monoclonal |
| Artikelnummer | BOT-BNCB0253-500 |
| Hersteller Artikelnummer | BNCB0253-500 |
| Alternativnummer | BOT-BNCB0253-500-500UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | FC, IHC, WB |
| Spezies Reaktivität | Bovine, Canine, Gallus, Human, Monkey, Mouse, Rabbit, Rat, Reptile |
| Immunogen | Human epidermal keratin |
| Konjugation | Biotin |
| Produktbeschreibung | This MAb recognizes the 56.5 kDa (CK10), 50 kDa (CK14), 50 kDa (CK15), 48 kDa (CK16), 40 kDa (CK19) keratins of the acidic (Type I or LMW) subfamily. Twenty human keratins are resolved with two-dimensional gel electrophoresis into acidic (pI 6.0) sub... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |
| Klon-Bezeichnung | [AE-1] |

| | |
|------------------------|--|
| Molekulargewicht | 40-56.5 kDa |
| UniProt | Q7Z794 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | <p>Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology formalin-fixed 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Western blotting 0.5-1 ug/mL Optimal dilution for a specific application should be determined by user</p> |