

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **IgM Immunoglobulin(IM260), Biotin conjugate, 0.1mg/mL, Clone: [IM260], Mouse, Monoclonal BOT-BNCB0260-500**

|                          |   |
|--------------------------|---|
| Artikelname              | IgM Immunoglobulin(IM260), Biotin conjugate, 0.1mg/mL, Clone: [IM260], Mouse, Monoclonal  |
| Artikelnummer            | BOT-BNCB0260-500  |
| Hersteller Artikelnummer | BNCB0260-500  |
| Alternativnummer         | BOT-BNCB0260-500-500UL  |
| Hersteller               | Biotium   |
| Wirt                     | Mouse   |
| Kategorie                | Antikörper  |
| Applikation              | IHC, WB   |
| Spezies Reaktivität      | Human   |
| Immunogen                | Recombinant heavy chain of human IgM  |
| Konjugation              | Biotin  |
| Produktbeschreibung      | Recognizes a protein of 75 kDa, identified as mu heavy chain of human immunoglobulins. It does not cross-react with alpha (IgA), gamma (IgG), epsilon (IgE), or delta (IgD), heavy chains, T-cells, monocytes, granulocytes, or erythrocytes. This MAb is ... |
| Klonalität               | Monoclonal  |
| Konzentration            | 0.1 mg/mL   |
| Klon-Bezeichnung         | [IM260]   |

|                        |   |
|------------------------|---|
| Molekulargewicht       | 50-75 kDa   |
| UniProt                | <a href="#">P01871</a>  |
| Puffer                 | PBS, 0.1% BSA, 0.05% azide  |
| Quelle                 | Animal  |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology formalin-fixed 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Western blotting 0.5-1 ug/mL Optimal dilution for a specific application should be determined by user |