

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Human IgM Immunoglobulin(ICO-30), Biotin conjugate, 0.1mg/mL, Clone: [ICO-30], Mouse, Monoclonal BOT-BNCB0364-100**

Artikelname	Human IgM Immunoglobulin(ICO-30), Biotin conjugate, 0.1mg/mL, Clone: [ICO-30], Mouse, Monoclonal
Artikelnummer	BOT-BNCB0364-100
Hersteller Artikelnummer	BNCB0364-100
Alternativnummer	BOT-BNCB0364-100-100UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Human
Immunogen	Heavy chain of human IgM
Konjugation	Biotin
Produktbeschreibung	Recognizes a protein of 75 kDa, identified as mu heavy chain of human immunoglobulins. It does not cross-react with alpha (IgA), gamma (IgG), epsilon (IgE), or delta (IgD), heavy chains, T-cells, monocytes, granulocytes, or erythrocytes. Monomeric Ig...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL
Klon-Bezeichnung	[ICO-30]

Molekulargewicht	50-75 kDa
UniProt	<a href="#">P01871</a>
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology formalin-fixed 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user