

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **CD6(C6/372), Biotin conjugate, 0.1mg/mL, IgG1, Clone: [C6/372], Mouse, Monoclonal BOT-BNCB0372-100**

|                          |   |
|--------------------------|---|
| Artikelname              | CD6(C6/372), Biotin conjugate, 0.1mg/mL, IgG1, Clone: [C6/372], Mouse, Monoclonal   |
| Artikelnummer            | BOT-BNCB0372-100  |
| Hersteller Artikelnummer | BNCB0372-100  |
| Alternativnummer         | BOT-BNCB0372-100-100UL  |
| Hersteller               | Biotium   |
| Wirt                     | Mouse   |
| Kategorie                | Antikörper  |
| Applikation              | IF, IHC   |
| Spezies Reaktivität      | Human   |
| Immunogen                | Human recombinant CD6 protein   |
| Konjugation              | Biotin  |
| Produktbeschreibung      | CD6 is a type I transmembrane glycoprotein that contains a 24-amino acid signal sequence, three extracellular scavenger receptor cysteine-rich (SRCR) domains, a membrane-spanning domain and a 44-amino acid cytoplasmic domain. The CD6 glycoprotein is ... |
| Klonalität               | Monoclonal  |
| Konzentration            | 0.1 mg/mL   |
| Klon-Bezeichnung         | [C6/372]  |

|                        |   |
|------------------------|---|
| Molekulargewicht       | 90-130 kDa  |
| Isotyp                 | IgG1  |
| UniProt                | <a href="#">P30203</a>  |
| Puffer                 | PBS, 0.1% BSA, 0.05% azide  |
| Quelle                 | Animal  |
| Anwendungsbeschreibung | <p>Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology formalin-fixed 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user</p> |