

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

p40(Rabbit PAb), Biotin conjugate, 0.1mg/mL, Clone: [PAb], Monoclonal BOT-BNCB0375-500

Artikelname	p40(Rabbit PAb), Biotin conjugate, 0.1mg/mL, Clone: [PAb], Monoclonal
Artikelnummer	BOT-BNCB0375-500
Hersteller Artikelnummer	BNCB0375-500
Alternativnummer	BOT-BNCB0375-500-500UL
Hersteller	Biotium
Wirt	Rabbit
Kategorie	Antikörper
Applikation	IHC, WB
Spezies Reaktivität	Bovine, Human, Mouse, Rat
Immunogen	A synthetic peptide (ENNAQTQFSEPQY) corresponding to aa5-17 of human p40
Konjugation	Biotin
Produktbeschreibung	p40 (p63 delta) is a marker determined to be highly specific for squamous basal cells by immunohistochemistry (IHC). The current more routinely recommended marker, p63, appears to have less specificity compared to p40, especially on squamous cell tum...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL
Klon-Bezeichnung	[PAb]

Molekulargewicht	40 kDa
UniProt	Q9H3D4
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology formalin-fixed 1-2 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 1 mM EDTA, pH 9.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Western blotting 0.5-1 ug/mL Optimal dilution for a specific application should be determined by user