

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Fascin-1(FSCN1/418), Biotin conjugate, 0.1mg/mL, Clone: [FSCN1/418], Mouse, Monoclonal BOT-BNCB0418-100**

|                          |   |
|--------------------------|---|
| Artikelname              | Fascin-1(FSCN1/418), Biotin conjugate, 0.1mg/mL, Clone: [FSCN1/418], Mouse, Monoclonal  |
| Artikelnummer            | BOT-BNCB0418-100  |
| Hersteller Artikelnummer | BNCB0418-100  |
| Alternativnummer         | BOT-BNCB0418-100-100UL  |
| Hersteller               | Biotium   |
| Wirt                     | Mouse   |
| Kategorie                | Antikörper  |
| Applikation              | FC, IF, IHC, WB   |
| Spezies Reaktivität      | Human, Rat  |
| Immunogen                | Full length recombinant human FSCN1 protein   |
| Konjugation              | Biotin  |
| Produktbeschreibung      | Recognizes a protein of 55 kDa, which is identified as fascin-1. Its actin binding ability is regulated by phosphorylation. Antibody to fascin-1 is a very sensitive marker for Reed-Sternberg cells and variants in nodular sclerosis, mixed cellularity,... |
| Klonalität               | Monoclonal  |
| Konzentration            | 0.1 mg/mL   |
| Klon-Bezeichnung         | [FSCN1/418]   |

|                        |   |
|------------------------|---|
| Molekulargewicht       | 55 kDa  |
| UniProt                | <a href="#">Q16658</a>  |
| Puffer                 | PBS, 0.1% BSA, 0.05% azide  |
| Quelle                 | Animal  |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology (formalin) Staining of formalin-fixed tissues requires boiling tissue sections in 1 mM EDTA, pH 8.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user |