

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **ODC-1 (Ornithine Decarboxylase-1)(ODC1/487), Biotin conjugate, 0.1mg/mL, Clone: [ODC1/487], Mouse, Monoclonal BOT-BNCB0487-500**

Artikelname	ODC-1 (Ornithine Decarboxylase-1)(ODC1/487), Biotin conjugate, 0.1mg/mL, Clone: [ODC1/487], Mouse, Monoclonal
Artikelnummer	BOT-BNCB0487-500
Hersteller Artikelnummer	BNCB0487-500
Alternativnummer	BOT-BNCB0487-500-500UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	FC, IF
Spezies Reaktivität	Human, Rat
Immunogen	Recombinant human ODC-1 protein
Konjugation	Biotin
Produktbeschreibung	Recognizes a 53 kDa protein, identified as the Ornithine Decarboxylase (ODC-1). ODC is the initial and rate-limiting enzyme in the biosynthetic pathway of polyamines and is involved in the conversion of ornithine to putrescine. The biological activit...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL
Klon-Bezeichnung	[ODC1/487]

Molekulargewicht	53 kDa
UniProt	<a href="#">P11926</a>
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology (formalin) Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Western blotting 0.5-1 ug/mL Optimal dilution for a specific application should be determined by user