

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Topoisomerase I, MT(TOP1MT/488), Biotin conjugate, 0.1mg/mL, Clone: [TOP1MT/488], Mouse, Monoclonal BOT-BNCB0488-100

| | |
|--------------------------|---|
| Artikelname | Topoisomerase I, MT(TOP1MT/488), Biotin conjugate, 0.1mg/mL, Clone: [TOP1MT/488], Mouse, Monoclonal |
| Artikelnummer | BOT-BNCB0488-100 |
| Hersteller Artikelnummer | BNCB0488-100 |
| Alternativnummer | BOT-BNCB0488-100-100UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | FC, IF, IHC, WB |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant full-length human TOP1MT protein |
| Konjugation | Biotin |
| Produktbeschreibung | DNA topoisomerases are nuclear enzymes that regulate the topological structure of DNA in eukaryotic cells by transiently breaking and rejoining DNA strands. Due to their roles in DNA replication, recombination, and transcription, DNA topoisomerases h... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |

| | |
|------------------------|--|
| Klon-Bezeichnung | [TOP1MT/488] |
| Molekulargewicht | 70 kDa |
| UniProt | Q969P6 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology formalin-fixed 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user |