

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Moesin(MSN/493), Biotin conjugate, 0.1mg/mL, Clone: [MSN/493], Mouse, Monoclonal BOT-BNCB0493-500**

Artikelname	Moesin(MSN/493), Biotin conjugate, 0.1mg/mL, Clone: [MSN/493], Mouse, Monoclonal
Artikelnummer	BOT-BNCB0493-500
Hersteller Artikelnummer	BNCB0493-500
Alternativnummer	BOT-BNCB0493-500-500UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	FC, IF, IHC, WB
Spezies Reaktivität	Human
Immunogen	Recombinant full-length human Moesin protein
Konjugation	Biotin
Produktbeschreibung	Recognizes 78 kDa moesin protein. Moesin, a member of the talin-4.1 superfamily, is a linking protein of the sub-membranous actin cytoskeleton. It is expressed in variable amounts in cells of different phenotypes such as macrophages, lymphocytes, fib...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL
Klon-Bezeichnung	[MSN/493]

Molekulargewicht	78 kDa
UniProt	<a href="#">P26038</a>
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Does not react with rat, others not known Immunohistology (formalin) Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Western blotting 0.5-1 ug/mL Optimal dilution for a specific application should be determined by user