

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Nucleolin(364-5), Biotin conjugate, 0.1mg/mL, Clone: [364-5], Mouse, Monoclonal BOT-BNCB0527-100

| | |
|--------------------------|---|
| Artikelname | Nucleolin(364-5), Biotin conjugate, 0.1mg/mL, Clone: [364-5], Mouse, Monoclonal |
| Artikelnummer | BOT-BNCB0527-100 |
| Hersteller Artikelnummer | BNCB0527-100 |
| Alternativnummer | BOT-BNCB0527-100-100UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | FC, IF, IHC, WB |
| Spezies Reaktivität | Human |
| Immunogen | Lysate of SU-DHL-1 Nuclei |
| Konjugation | Biotin |
| Produktbeschreibung | Recognizes Nucleolin (NCL), the major nucleolar phosphoprotein of growing eukaryotic cells. NCL is located mainly in dense fibrillar regions of the nucleolus. It is found associated with intranucleolar chromatin and pre-ribosomal particles. Human NCL... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |
| Klon-Bezeichnung | [364-5] |

| | |
|------------------------|--|
| Molekulargewicht | 76 kDa (predicted), 100-110 kDa (apparent) |
| UniProt | P19338 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Does not react with mouse, rat or cow, others not known Immunohistology formalin-fixed 0.25-0.5 ug/mL Staining of formalin-fixed tissues is enhanced by boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Western blotting 0.5-1 ug/mL Optimal dilution for a specific application should be determined by user |