

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **CD45 / LCA(2B11 + PD7/26), Biotin conjugate, 0.1mg/mL, Clone: [2B11 PD7/26], Mouse, Monoclonal BOT-BNCB0745-100**

|                          |   |
|--------------------------|---|
| Artikelname              | CD45 / LCA(2B11 + PD7/26), Biotin conjugate, 0.1mg/mL, Clone: [2B11 PD7/26], Mouse, Monoclonal  |
| Artikelnummer            | BOT-BNCB0745-100  |
| Hersteller Artikelnummer | BNCB0745-100  |
| Alternativnummer         | BOT-BNCB0745-100-100UL  |
| Hersteller               | Biotium   |
| Wirt                     | Mouse   |
| Kategorie                | Antikörper  |
| Applikation              | FC, IF, IHC, IP   |
| Spezies Reaktivität      | Canine, Human   |
| Immunogen                | Isolated neoplastic cells from T cell lymphoma (2B11), and human peripheral blood lymphocytes maintained in T cell growth factor (PD7/26)   |
| Konjugation              | Biotin  |
| Produktbeschreibung      | CD45R, also designated CD45 and PTPRC, has been identified as a transmembrane glycoprotein, broadly expressed among hematopoietic cells. Multiple isoforms of CD45R are distributed throughout the immune system according to cell type. These isoforms ar... |
| Klonalität               | Monoclonal  |

|                        |  |
|------------------------|--|
| Konzentration          | 0.1 mg/mL  |
| Klon-Bezeichnung       | [2B11 PD7/26]  |
| Molekulargewicht       | 180-220 kDa  |
| UniProt                | <a href="#">P08575</a>   |
| Puffer                 | PBS, 0.1% BSA, 0.05% azide   |
| Quelle                 | Animal   |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology (formalin) Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user |