

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Cytokeratin 7(KRT7/760), Biotin conjugate, 0.1mg/mL, IgG1, Clone: [KRT7/760], Mouse, Monoclonal BOT-BNCB0760-100

| | |
|--------------------------|---|
| Artikelname | Cytokeratin 7(KRT7/760), Biotin conjugate, 0.1mg/mL, IgG1, Clone: [KRT7/760], Mouse, Monoclonal |
| Artikelnummer | BOT-BNCB0760-100 |
| Hersteller Artikelnummer | BNCB0760-100 |
| Alternativnummer | BOT-BNCB0760-100-100UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | FC, IF, IHC |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant full-length human KRT7 protein |
| Konjugation | Biotin |
| Produktbeschreibung | This antibody recognizes an intermediate filament protein (IFP) of 55 kDa, which is identified as cytokeratin 7. This MAb is highly specific to cytokeratin 7 and shows no cross-reaction with other IFPs. Cytokeratin 7 is a basic cytokeratin, which is ... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |
| Klon-Bezeichnung | [KRT7/760] |

| | |
|------------------------|---|
| Molekulargewicht | 55 kDa |
| Isotyp | IgG1 |
| UniProt | P08729 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | <p>Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology (formalin) Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user</p> |