

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

IgM Immunoglobulin(Cocktail), Biotin conjugate, 0.1mg/mL, Clone: [Cocktail], Mouse, Monoclonal BOT-BNCB0762-100

Artikelname	IgM Immunoglobulin(Cocktail), Biotin conjugate, 0.1mg/mL, Clone: [Cocktail], Mouse, Monoclonal
Artikelnummer	BOT-BNCB0762-100
Hersteller Artikelnummer	BNCB0762-100
Alternativnummer	BOT-BNCB0762-100-100UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Human
Immunogen	Recombinant heavy chain of human IgM (IM260 & IM373), Heavy chain of human IgM (ICO-30)
Konjugation	Biotin
Produktbeschreibung	Recognizes a protein of 75 kDa, identified as mu heavy chain of human immunoglobulins. It does not cross-react with alpha (IgA), gamma (IgG), epsilon (IgE), or delta (IgD), heavy chains, T-cells, monocytes, granulocytes, or erythrocytes. This MAb is ...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL

Klon-Bezeichnung	[Cocktail]
Molekulargewicht	50-75 kDa
UniProt	P01871
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology formalin-fixed 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Western blotting 0.5-1 ug/mL Optimal dilution for a specific application should be determined by user