

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

CD20 / MS4A1 (B-Cell Marker)(L26), Biotin conjugate, 0.1mg/mL, Clone: [L26], Mouse, Monoclonal BOT-BNCB0766-100

Artikelname	CD20 / MS4A1 (B-Cell Marker)(L26), Biotin conjugate, 0.1mg/mL, Clone: [L26], Mouse, Monoclonal
Artikelnummer	BOT-BNCB0766-100
Hersteller Artikelnummer	BNCB0766-100
Alternativnummer	BOT-BNCB0766-100-100UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	FC, IF, IHC, WB
Spezies Reaktivität	Human
Immunogen	Human tonsil B cells
Konjugation	Biotin
Produktbeschreibung	This antibody recognizes a protein of 30-33 kDa, which is identified as CD20. Its epitope is located in the cytoplasmic domain of CD20 and was, therefore, ascribed as CD20cy in the 5th Workshop. CD20 is a non-Ig differentiation antigen of B-cells and...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL
Klon-Bezeichnung	[L26]

Molekulargewicht	33-37 kDa
UniProt	P11836
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistochemistry (formalin-fixed): 0.5-1 ug/mL for 30 minutes at RT Western blot: 1-2 ug/mL Flow cytometry: 0.5-1 ug/million cells Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user