

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **CD20 / MS4A1 (B-Cell Marker)(L26), Biotin conjugate, 0.1mg/mL, Clone: [L26], Mouse, Monoclonal BOT-BNCB0766-500**

|                          |   |
|--------------------------|---|
| Artikelname              | CD20 / MS4A1 (B-Cell Marker)(L26), Biotin conjugate, 0.1mg/mL, Clone: [L26], Mouse, Monoclonal  |
| Artikelnummer            | BOT-BNCB0766-500  |
| Hersteller Artikelnummer | BNCB0766-500  |
| Alternativnummer         | BOT-BNCB0766-500-500UL  |
| Hersteller               | Biotium   |
| Wirt                     | Mouse   |
| Kategorie                | Antikörper  |
| Applikation              | FC, IF, IHC, WB   |
| Spezies Reaktivität      | Human   |
| Immunogen                | Human tonsil B cells  |
| Konjugation              | Biotin  |
| Produktbeschreibung      | This antibody recognizes a protein of 30-33 kDa, which is identified as CD20. Its epitope is located in the cytoplasmic domain of CD20 and was, therefore, ascribed as CD20cy in the 5th Workshop. CD20 is a non-Ig differentiation antigen of B-cells and... |
| Klonalität               | Monoclonal  |
| Konzentration            | 0.1 mg/mL   |
| Klon-Bezeichnung         | [L26]   |

|                        |   |
|------------------------|---|
| Molekulargewicht       | 33-37 kDa   |
| UniProt                | <a href="#">P11836</a>  |
| Puffer                 | PBS, 0.1% BSA, 0.05% azide  |
| Quelle                 | Animal  |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistochemistry (formalin-fixed): 0.5-1 ug/mL for 30 minutes at RT Western blot: 1-2 ug/mL Flow cytometry: 0.5-1 ug/million cells Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user |