

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Cytokeratin 17(KRT17/778), Biotin conjugate, 0.1mg/mL, Clone: [KRT17/778], Mouse, Monoclonal BOT-BNCB0778-100

Artikelname	Cytokeratin 17(KRT17/778), Biotin conjugate, 0.1mg/mL, Clone: [KRT17/778], Mouse, Monoclonal
Artikelnummer	BOT-BNCB0778-100
Hersteller Artikelnummer	BNCB0778-100
Alternativnummer	BOT-BNCB0778-100-100UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Bovine, Goat, Human, Porcine, Rat
Immunogen	Recombinant full-length human KRT17 protein
Konjugation	Biotin
Produktbeschreibung	Cytokeratin 17 (CK17) is normally expressed in the basal cells of complex epithelia but not in stratified or simple epithelia. Antibody to CK17 is an excellent tool to distinguish myoepithelial cells from luminal epithelium of various glands such as ...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL
Klon-Bezeichnung	[KRT17/778]

Molekulargewicht	46 kDa
UniProt	Q04695
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology formalin-fixed 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user