

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

bcl-2(BCL2/782), Biotin conjugate, 0.1mg/mL, Clone: [BCL2/782], Mouse, Monoclonal BOT-BNCB0782-100

| | |
|--------------------------|---|
| Artikelname | bcl-2(BCL2/782), Biotin conjugate, 0.1mg/mL, Clone: [BCL2/782], Mouse, Monoclonal |
| Artikelnummer | BOT-BNCB0782-100 |
| Hersteller Artikelnummer | BNCB0782-100 |
| Alternativnummer | BOT-BNCB0782-100-100UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC, WB |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant full-length human bcl-2 protein |
| Konjugation | Biotin |
| Produktbeschreibung | This antibody recognizes a protein of 25-26 kDa, identified as Bcl-2. This antibody shows no cross-reaction with Bcl-x or Bax proteins. Bcl-2 suppresses apoptosis in a variety of cell systems including factor-dependent lymphohematopoietic and neural ... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |
| Klon-Bezeichnung | [BCL2/782] |

| | |
|------------------------|--|
| Molekulargewicht | 25-26 kDa |
| UniProt | P10415 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology formalin-fixed 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM Tris with 1 mM EDTA, pH 9.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Western blotting 0.5-1 ug/mL Optimal dilution for a specific application should be determined by user |