

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

CD30 / TNFRSF8 (Hodgkin & Reed-Sternberg Cell Marker)(Ber-H2 + CD30/412), Biotin conjugate, 0.1mg/mL, Clone: [Ber-H2 CD30/412], Mouse, Monoclonal BOT-BNCB0849-500

Artikelname	CD30 / TNFRSF8 (Hodgkin & Reed-Sternberg Cell Marker)(Ber-H2 + CD30/412), Biotin conjugate, 0.1mg/mL, Clone: [Ber-H2 CD30/412], Mouse, Monoclonal
Artikelnummer	BOT-BNCB0849-500
Hersteller Artikelnummer	BNCB0849-500
Alternativnummer	BOT-BNCB0849-500-500UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Spezies Reaktivität	Human
Immunogen	Co cell line established from a patient with Hodgkin's disease of T-cell lineage (Ber-H2), human CD30 recombinant protein (CD30/412)
Konjugation	Biotin
Produktbeschreibung	This antibody recognizes a single chain glycoprotein of 105/120 kDa, identified as CD30/Ki-1. CD30 is synthesized as a 90 kDa precursor, which is processed in the Golgi complex into a membrane-bound phosphorylated mature 105/120 kDa glycoprotein. In ...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL

Klon-Bezeichnung	[Ber-H2 CD30/412]
Molekulargewicht	105-120 kDa
UniProt	P28908
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	For coating for ELISA, order Ab without BSA Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Optimal dilution and staining procedure for a specific application should be determined by user Recommended starting concentrations for titration are 1-2 ug/mL for most applications, or 1 ug/million cells/100 uL for flow cytometry