

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

CD20 / MS4A1 (B-Cell Marker)(L26 + IGEL/773), Biotin conjugate, 0.1mg/mL, Clone: [L26 IGEL/773], Mouse, Monoclonal BOT-BNCB0854-100

Artikelname	CD20 / MS4A1 (B-Cell Marker)(L26 + IGEL/773), Biotin conjugate, 0.1mg/mL, Clone: [L26 IGEL/773], Mouse, Monoclonal
Artikelnummer	BOT-BNCB0854-100
Hersteller Artikelnummer	BNCB0854-100
Alternativnummer	BOT-BNCB0854-100-100UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Spezies Reaktivität	Human, Monkey, Primate
Immunogen	Human tonsil B cells (L26), Recombinant human MS4A1 protein (IGEL/773)
Konjugation	Biotin
Produktbeschreibung	This antibody recognizes a protein of 30-33 kDa, which is identified as CD20. Its epitope is located in the cytoplasmic domain of CD20 and was, therefore, ascribed as CD20cy in the 5th Workshop. CD20 is a non-Ig differentiation antigen of B-cells and...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL
Klon-Bezeichnung	[L26 IGEL/773]

Molekulargewicht	33-37 kDa
UniProt	P11836
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Does not react with Cow, Dog, Pig or Rat For coating for ELISA, order Ab without BSA Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Optimal dilution and staining procedure for a specific application should be determined by user Recommended starting concentrations for titration are 1-2 ug/mL for most applications, or 1 ug/million cells/100 uL for flow cytometry