

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **WT1 (Wilms Tumor 1)(WT1/857), Biotin conjugate, 0.1mg/mL, Clone: [WT1/857], Mouse, Monoclonal BOT-BNCB0857-500**

Artikelname	WT1 (Wilms Tumor 1)(WT1/857), Biotin conjugate, 0.1mg/mL, Clone: [WT1/857], Mouse, Monoclonal
Artikelnummer	BOT-BNCB0857-500
Hersteller Artikelnummer	BNCB0857-500
Alternativnummer	BOT-BNCB0857-500-500UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Human, Mouse, Rat
Immunogen	Recombinant human WT1 protein
Konjugation	Biotin
Produktbeschreibung	Recognizes a 47-55 kDa-tumor suppressor protein, identified as Wilms Tumor (WT1) protein. The antibody reacts with all isoforms of the full-length WT1 and also identifies WT1 lacking exon 2-encoded amino acids, frequently found in subsets of sporadic...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL
Klon-Bezeichnung	[WT1/857]

Molekulargewicht	47-55 kDa
UniProt	<a href="#">P19544</a>
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology formalin-fixed 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user