

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **p34 / cdk1(CDK1/873), Biotin conjugate, 0.1mg/mL, Clone: [CDK1/873], Mouse, Monoclonal BOT-BNCB0873-500**

Artikelname	p34 / cdk1(CDK1/873), Biotin conjugate, 0.1mg/mL, Clone: [CDK1/873], Mouse, Monoclonal
Artikelnummer	BOT-BNCB0873-500
Hersteller Artikelnummer	BNCB0873-500
Alternativnummer	BOT-BNCB0873-500-500UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC, WB
Spezies Reaktivität	Bovine, Human, Mink, Monkey
Immunogen	Recombinant human CDK1 protein
Konjugation	Biotin
Produktbeschreibung	Recognizes a 34 kDa protein, which is identified as cyclin dependent kinase 1 (cdk1) or p34cdc2 protein kinase. cdk1 / p34cdc2 plays a crucial role during cell division and is most active during mitosis. It is predominantly localized in the nucleus. ...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL
Klon-Bezeichnung	[CDK1/873]

Molekulargewicht	34 kDa
UniProt	<a href="#">P06493</a>
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Does not react with mouse, rat, Xenopus, Drosophila, S. pombe, or S. cerevisiae, others not known Immunohistology formalin-fixed 2-4 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user