

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Human IgG Immunoglobulin(B33/20), Biotin conjugate, 0.1mg/mL, Clone: [B33/20], Mouse, Monoclonal BOT-BNCB0952-100

Artikelname	Human IgG Immunoglobulin(B33/20), Biotin conjugate, 0.1mg/mL, Clone: [B33/20], Mouse, Monoclonal
Artikelnummer	BOT-BNCB0952-100
Hersteller Artikelnummer	BNCB0952-100
Alternativnummer	BOT-BNCB0952-100-100UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Human
Immunogen	Purified polyclonal human Ig Gamma Chain
Konjugation	Biotin
Produktbeschreibung	Recognizes a protein of 75 kDa, identified as i heavy chain of human immunoglobulins. Its epitope maps in CH2 domain of Fc region of IgG. It reacts with all sub-classes of i chain of human immunoglobulins. It does not cross-react with i (IgA), i (IgM...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL
Klon-Bezeichnung	[B33/20]

Molekulargewicht	75 kDa
UniProt	P01857
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology formalin/paraffin 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user